

WARNING | TOXIC ECOTOXIC
KEEP OUT OF REACH OF CHILDREN

ExtractaTM
Herbicides

Performa Tribenuron-Methyl 750WG

Active Ingredient: 750g/kg Tribenuron-methyl
in the form of a water dispersible granule

GROUP 2 HERBICIDE

PerformaTM
Crop Protection

A selective herbicide for control of certain
broadleaf weeds in WHEAT, BARLEY and OATS,
and in CONSERVATION TILLAGE PROGRAMS



READ SAFETY DIRECTIONS BEFORE OPENING OR USING

Registered pursuant to the ACVM Act 1997, No. P010484
See www.foodsafety.govt.nz for registration conditions.
Approved pursuant to the HSNO Act 1996, No. HSR000234
See www.epa.govt.nz for approval controls.

Batch No:

Date of Manufacture:

Net Contents:

2 KG

READ LABEL COMPLETELY BEFORE OPENING AND USING PRODUCT

GHS CLASSIFICATION

Eye irritation Category 2, Skin sensitization Category 1, Specific target organ toxicity (repeated exposure) Category 2, Hazardous to soil organisms, Hazardous to the aquatic environment acute Category 1, Hazardous to the aquatic environment chronic Category 1.

TOXICITY

WARNING: Causes serious eye irritation. May cause an allergic skin reaction. May cause organ damage from repeated or prolonged oral exposure.

ECOTOXICITY

Very toxic to aquatic life and with long lasting effects. Hazardous to soil organisms. **Selective Herbicide – Very toxic to some plant species.** Some plants may be damaged or killed by contact with this product. Do not apply directly onto or into water. Use this product carefully. Do not contaminate aquatic environments with product, spray drift or empty container. Collect spillage. Spray drift or equipment contamination may cause serious damage to desirable plants. Do not allow spray drift outside the target area to occur. Do not use or flush equipment near to desirable plants or their roots. Do not use the same equipment to apply other chemicals to susceptible crops unless thoroughly cleaned and decontaminated.



PERSONNEL QUALIFICATION & EXPERIENCE:

This product must be under the control of a suitably qualified person when applied into or onto water, and the water has the potential to leave the application area. For a list of suitable qualifications refer to the Hazardous Substances (Hazardous Property Controls) Notice 2017 Subpart C. This product must not be used for any purpose, or in any manner, contrary to this label unless authorised under appropriate legislation.

RECORD KEEPING: Records must be kept whenever 3kg or more of Performa Tribenuron-Methyl 750WG is applied within 24 hours, in a place where the substance is likely to enter air or water and leave the application area, a PCBU with management or control of the substance must ensure that a written record is kept of each application of the substance.

PRECAUTIONS

PROTECTIVE EQUIPMENT: Avoid contact with eyes and skin. Avoid inhalation of dust or spray mist. When mixing or applying, wear overalls, boots, gloves and eye protection. Clean protective clothing daily after work. Do not eat, drink or smoke while using. Remove protective clothing and wash hands and face thoroughly before meals and after work.

STORAGE: Keep out of reach of children. Store in the original tightly closed container in a secure area. When stored appropriately this product should show no significant degradation for two years from the date of manufacture. Contact your supplier for further information about the use of product that is older than this.

DISPOSAL: Dispose of this product by using it in accordance with this label. Dispose of the container via AgRecovery after triple rinsing (use the rinsate for spraying). Do not burn. Do not use the container for any other purpose.

EQUIPMENT: Apply with well-maintained and calibrated spray equipment.

First Aid: If swallowed, Rinse mouth. Do NOT induce vomiting. Get medical advice/attention if you feel unwell. If skin or hair contact occurs, remove contaminated clothing and flush skin and hair with running water. If skin irritation or rash occurs: Get medical advice/attention. If splashed in eyes, wash out immediately with water. If eye irritation persists: Get medical advice/attention. If inhaled: Remove to fresh air and keep at rest in a comfortable position for breathing. For advice, contact the National Poisons Centre 0800 POISON (0800 764 766) or a doctor. If medical advice is needed, have product container or label at hand.

Emergency Response: In a transport emergency call 111, Police or Fire Brigade. For specialist advice in an emergency only, call 0800 CHEMCALL (0800 243 622) (24 hrs).

Spillage: Exclude non-essential people from the area. Wear protective clothing and prevent material from entering waterways. Absorb spill with inert material such as soil, sand, etc, and place in waste container. Wash area with water, and absorb with further inert material. Dispose of safely to a suitable landfill.

Safety Data Sheet: Additional information is provided in the safety data sheet which is available from the distributor.

SPRAYER DECONTAMINATION: Always wash tank and flush equipment thoroughly with clean water after use. To avoid subsequent injury to sensitive crops, immediately after spraying thoroughly remove all traces of Performa Tribenuron-Methyl 750WG from mixing and spraying equipment as follows:

1. Drain tank, then flush tank, booms and hoses with clean water for a minimum of 10 minutes.
2. Fill the tank with clean water and add 1litre chlorine bleach (containing 3% solution sodium hypochlorite) per 100L of water. Flush through booms with agitation then drain.
3. Repeat step 2.
4. Nozzles and filters should be removed and cleaned separately.
5. To remove traces of chlorine bleach, rinse tank thoroughly with clean water and flush through hoses and boom

WARRANTY: This formulation is warranted to contain the specified active ingredient within accepted analytical tolerance when packed. The user bears the risk for damage resulting from factors beyond the manufacturer's control. The manufacturer declines all liability for damage resulting from improper storage or use of the product.

GENERAL INFORMATION

PERFORMA TRIBENURON-METHYL 750WG is a sulfonylurea herbicide which is non-hormonal and non-volatile. Performa Tribenuron-Methyl 750WG works best when applied to weeds that are small and actively growing. Susceptible weeds will stop growing soon after treatment but symptoms may take 1-3 weeks to appear. Performa Tribenuron-Methyl 750WG works quicker under warm, moist growth conditions and slower under cold, dry conditions following application. Do NOT spray crops that are under stress from drought, water logging, disease or insect damage, as crop damage may occur. Crop damage may also occur if these conditions occur after spraying. Do NOT apply Performa Tribenuron-Methyl 750WG to crops under-sown with white clover or lucerne. Take all reasonable steps to ensure that the substance does not cause any significant adverse effects to the environment beyond the application area.

MIXING: Add the required quantity of Performa Tribenuron-Methyl 750WG to spray tank partly filled with water and maintain agitation through filling and spraying. Ensure Performa Tribenuron-Methyl 750WG is thoroughly mixed before adding any other pesticide to the tank. Use all spray solution within 24hr as degradation may occur.

COMPATIBILITY: Performa Tribenuron-Methyl 750WG is compatible with most commonly used herbicides, insecticides and fungicides.

RAINFALL: Rainfall within 1 hour of application may reduce the effectiveness of Performa Tribenuron-Methyl 750WG.

RESISTANCE MANAGEMENT:

GROUP 2 HERBICIDE

Performa Tribenuron-Methyl 750WG contains Tribenuron-methyl, a herbicide from the sulfonylurea chemical group with the inhibition of acetolactate synthase ALS mode of action. Naturally occurring weed biotypes resistant to sulfonylurea herbicides are known to exist. These become more prevalent after many years of continuous use of the product. To prevent or delay development of resistant biotypes, use Performa Tribenuron-Methyl 750WG in tank mixture or rotation with herbicides of a different mode of action against the weed. Consult www.resistance.nzpps.org/ for alternative modes of action and details of resistance management strategies (if any) for the crops listed on the label.

DIRECTIONS FOR USE: Do NOT spray crops that are under stress from drought, water saturated soil, disease or insect damage, as crop damage may occur. Crop damage may also occur if these conditions occur after spraying. Do NOT apply to crops undersown with clover or lucerne. Use a minimum of 60 litres of water per hectare in all methods of application. Crop injury may result from spray overlap. When starting, slowing or stopping, shut off the spray boom.

It is an offence for users of this product to cause residues exceeding the relevant MRL in the Food Notice: Maximum Residue Levels for Agricultural Compounds.

WITHHOLDING PERIOD: Not required when used as directed.

CROP / SITUATION	RATE	WEEDS CONTROLLED	CRITICAL COMMENTS
Cereals – Wheat, Barley, Oats	15g/ha	Californian thistle, chickweed, fathen, scarlet pimpernel, scotch thistle, shepherds purse, spurry, willow weed. <i>*Apply to rosettes (10 – 15cm tall); thistles emerging after application will not be controlled.</i>	Apply at any stage from the 2-leaf stage up to just before the 'boot' stage. For best results spray before the crop canopy closes to small actively growing weeds. Use a minimum of 60 litres/ha of water. When used alone, add a non-ionic surfactant at label recommended rate.
	25g/ha	Redroot, twin cross, wireweed, sheep's sorrel.	The higher rate is required for these weeds.
Barley only	15g/ha		To improve the spectrum of weeds controlled, follow the label recommendations of the tank mix partner.
Conservation Tillage – Cereals, New Pasture, Peas, Potatoes, Squash	40g/ha	Clovers, ragwort, <i>yarrow 'Ragwort plants may be more palatable after spraying, do NOT graze until treated plants brown out.</i>	For improved control of certain broadleaf weeds, use in tank mix with glyphosate plus penetrant before the crop is sown. Plantback Period: Do not sow the crop within 14 days of using PERFORMA TRIBENURON-METHYL 750WG. If sown 14 – 28 days after application, some brassica cultivars may suffer loss of vigour during early establishment. However, this has no longer term effects. Do not sow if the soil pH exceeds 6.5. Do not use PERFORMA TRIBENURON-METHYL 750WG within 2 months after applying lime.
Conservation Tillage – Brassicas	40g/ha		For improved control of certain broadleaf weeds, use in tank mix with glyphosate plus penetrant before the crop is sown. Plantback Period: Do not sow the crop within 14 days of using PERFORMA TRIBENURON-METHYL 750WG. If sown 14 – 28 days after application, some brassica cultivars may suffer loss of vigour during early establishment. However, this has no longer term effects. Do not sow if the soil pH exceeds 6.5. Do not use PERFORMA TRIBENURON-METHYL 750WG within 2 months after applying lime.

MANUFACTURED FOR
WHOLESALE SEEDS LTD BY:

Danken New Zealand Limited.
Telephone: 021 241 6152